

Cat. No. 64-2313B

## **QUICK-SETTING EPOXY**

### **DIRECTIONS**

1. Wipe surfaces clean and dry where you plan to apply the adhesive.
2. Cut both nozzles at the tip.
3. Using the plastic blister packaging as a mixing tray, press down slowly on the plunger to dispense adhesive and resin. Mix thoroughly.
4. Use a toothpick to apply the mixture to both surfaces to be bonded. Then use a clamp, tape, or weights to hold them together for at least 4-6 minutes.
  - Do not handle the item for about 30 minutes after applying the adhesive.
  - The bond reaches its full strength after 16 hours.
5. To reseal the container, remove the cap from between the syringes, pull the plunger back slightly, clean the nozzles, and place the cap over the nozzles.

**Note:** the cap only fits one way.

**WARNING!** KEEP OUT OF REACH OF CHILDREN. USE ONLY AS DIRECTED. INTENTIONAL MISUSE BY INHALING, INGESTING OR EATING THE CONTENTS CAN BE HARMFUL OR CAUSE DEATH.

**CAUTION!** May cause skin or eye irritation. Avoid prolonged breathing of vapors and contact with eyes and skin. For Medical Emergencies call (303) 623-5716

Contents: Contenta Braphenol A/Epichlorohydnn based epoxy resin CAS 25068.38.6. Proprietary Mercaotan Polymer.

**FIRST AID: EYES:** In case of eye contact, flush with water for 15 minutes and get immediate medical attention. **SKIN:** Wash with soap and water for 15 minutes. Wash ccontaminated clothing before reuse. **INGESTION:** DO NOT induce vomiting. Seek immediate medical attention. **INHALATION:** Remove to fresh air. Remove to fresh air. Adminster oxygen if breathing is difficult. Seek immediate medical attention.